

AD _____

GRANT NUMBER: DAMD17-94-J-4050


TITLE: General Methods for Identifying G1 Substrates of the Cdk
Protein Kinases

PRINCIPAL INVESTIGATOR: Dr. A. B. Futcher

CONTRACTING ORGANIZATION: Cold Spring Harbor Laboratory
Cold Spring Harbor, New York 11724

REPORT DATE: June 27, 1995

TYPE OF REPORT: Annual



PREPARED FOR: U.S. Army Medical Research and Materiel Command
Fort Detrick, Maryland 21702-5012

DISTRIBUTION STATEMENT: Approved for public release;
distribution unlimited

The views, opinions and/or findings contained in this report are those of the author(s) and should not be construed as an official Department of the Army position, policy or decision unless so designated by other documentation.

19950901 044

DTIC QUALITY INSPECTED 8

REPORT DOCUMENTATION PAGE			Form Approved OMB No. 0704-0188	
Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503.				
1. AGENCY USE ONLY (Leave blank)		2. REPORT DATE June 27, 1995	3. REPORT TYPE AND DATES COVERED Annual 1 Jun 94 - 31 May 95	
4. TITLE AND SUBTITLE General Methods for Identifying G1 - Phase Substrates of Cdk Cyclin Complexes			5. FUNDING NUMBERS DAMD17-94-J-4050	
6. AUTHOR(S) Dr. A. B. Futcher				
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Cold Spring Harbor Laboratory Cold Spring Harbor, New York 11724			8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES) U.S. Army Medical Research and Materiel Command Fort Detrick, Maryland 21702-5012			10. SPONSORING/MONITORING AGENCY REPORT NUMBER	
11. SUPPLEMENTARY NOTES				
12a. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release; distribution unlimited			12b. DISTRIBUTION CODE	
13. ABSTRACT (Maximum 200 words) We are using two-dimensional gels in combination with genetic and biochemical techniques to identify G1-phase substrates of cyclin-Cdk complexes. We have optimized protocols for labelling proteins with ³² P prior to two-dimensional gel electrophoresis. Yeast strains have been constructed bearing six different cyclins expressed from a GAL promoter, to allow inducible phosphorylation of the desired substrates. A commercial antibody has been identified and obtained which may specifically immunoprecipitate phosphorylated forms of the desired substrates. Putative substrates have been examined <i>in vitro</i> . Together, these accomplishments provide a good platform from which to identify real, <i>in vivo</i> substrates in the near future.				
14. SUBJECT TERMS Cdks/Cyclins/Cdk Substrates			15. NUMBER OF PAGES 7	
			16. PRICE CODE	
17. SECURITY CLASSIFICATION OF REPORT Unclassified	18. SECURITY CLASSIFICATION OF THIS PAGE Unclassified	19. SECURITY CLASSIFICATION OF ABSTRACT Unclassified	20. LIMITATION OF ABSTRACT Unlmtied	

GENERAL INSTRUCTIONS FOR COMPLETING SF 298

The Report Documentation Page (RDP) is used in announcing and cataloging reports. It is important that this information be consistent with the rest of the report, particularly the cover and title page. Instructions for filling in each block of the form follow. It is important to **stay within the lines** to meet **optical scanning requirements**.

Block 1. Agency Use Only (Leave blank).

Block 2. Report Date. Full publication date including day, month, and year, if available (e.g. 1 Jan 88). Must cite at least the year.

Block 3. Type of Report and Dates Covered. State whether report is interim, final, etc. If applicable, enter inclusive report dates (e.g. 10 Jun 87 - 30 Jun 88).

Block 4. Title and Subtitle. A title is taken from the part of the report that provides the most meaningful and complete information. When a report is prepared in more than one volume, repeat the primary title, add volume number, and include subtitle for the specific volume. On classified documents enter the title classification in parentheses.

Block 5. Funding Numbers. To include contract and grant numbers; may include program element number(s), project number(s), task number(s), and work unit number(s). Use the following labels:

C - Contract	PR - Project
G - Grant	TA - Task
PE - Program Element	WU - Work Unit Accession No.

Block 6. Author(s). Name(s) of person(s) responsible for writing the report, performing the research, or credited with the content of the report. If editor or compiler, this should follow the name(s).

Block 7. Performing Organization Name(s) and Address(es). Self-explanatory.

Block 8. Performing Organization Report Number. Enter the unique alphanumeric report number(s) assigned by the organization performing the report.

Block 9. Sponsoring/Monitoring Agency Name(s) and Address(es). Self-explanatory.

Block 10. Sponsoring/Monitoring Agency Report Number. (If known)

Block 11. Supplementary Notes. Enter information not included elsewhere such as: Prepared in cooperation with...; Trans. of...; To be published in.... When a report is revised, include a statement whether the new report supersedes or supplements the older report.

Block 12a. Distribution/Availability Statement. Denotes public availability or limitations. Cite any availability to the public. Enter additional limitations or special markings in all capitals (e.g. NOFORN, REL, ITAR).

DOD - See DoDD 5230.24, "Distribution Statements on Technical Documents."

DOE - See authorities.

NASA - See Handbook NHB 2200.2.

NTIS - Leave blank.

Block 12b. Distribution Code.

DOD - Leave blank.

DOE - Enter DOE distribution categories from the Standard Distribution for Unclassified Scientific and Technical Reports.

NASA - Leave blank.

NTIS - Leave blank.

Block 13. Abstract. Include a brief (*Maximum 200 words*) factual summary of the most significant information contained in the report.

Block 14. Subject Terms. Keywords or phrases identifying major subjects in the report.

Block 15. Number of Pages. Enter the total number of pages.

Block 16. Price Code. Enter appropriate price code (*NTIS only*).

Blocks 17. - 19. Security Classifications. Self-explanatory. Enter U.S. Security Classification in accordance with U.S. Security Regulations (i.e., UNCLASSIFIED). If form contains classified information, stamp classification on the top and bottom of the page.

Block 20. Limitation of Abstract. This block must be completed to assign a limitation to the abstract. Enter either UL (unlimited) or SAR (same as report). An entry in this block is necessary if the abstract is to be limited. If blank, the abstract is assumed to be unlimited.

FOREWORD

Opinions, interpretations, conclusions and recommendations are those of the author and are not necessarily endorsed by the US Army.

NA Where copyrighted material is quoted, permission has been obtained to use such material.

NA Where material from documents designated for limited distribution is quoted, permission has been obtained to use the material.

B7 Yes Citations of commercial organizations and trade names in this report do not constitute an official Department of Army endorsement or approval of the products or services of these organizations.

B7 Yes In conducting research using animals, the investigator(s) adhered to the "Guide for the Care and Use of Laboratory Animals," prepared by the Committee on Care and Use of Laboratory Animals of the Institute of Laboratory Resources, National Research Council (NIH Publication No. 86-23, Revised 1985).

NA For the protection of human subjects, the investigator(s) adhered to policies of applicable Federal Law 45 CFR 46.

B7 Yes In conducting research utilizing recombinant DNA technology, the investigator(s) adhered to current guidelines promulgated by the National Institutes of Health.

B7 Yes In the conduct of research utilizing recombinant DNA, the investigator(s) adhered to the NIH Guidelines for Research Involving Recombinant DNA Molecules.

B7 Yes In the conduct of research involving hazardous organisms, the investigator(s) adhered to the CDC-NIH Guide for Biosafety in Microbiological and Biomedical Laboratories.

Bruce Futch June 23, 1995
PI - Signature Date

N.A. = Not Applicable.

TABLE OF CONTENTS

Front Cover	1
SF 298 - Report Documentation Page	2
Foreword	3
Table of Contents	4
Introduction	5
Body of Report	5
Conclusions	7
References	7

Accession For	
DTIC GRA&I	<input checked="checked" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By	
Distribution/	
Availability Codes	
Dist	Avail and/or Special
A-1	

5. Introduction.

There is general agreement that in all eukaryotes, phosphorylation by various cyclin-Cdk complexes controls and orchestrates key cell cycle events. These events include commitment in G1 phase, initiation of DNA synthesis in S phase, and spindle formation and elongation in mitosis. However, despite knowing a great deal about the cyclin-Cdk complexes themselves, and despite years of investigation by many laboratories, we know only about half a dozen substrates of the cyclin-Cdk kinases, and none of these explain the control of critical cell cycle events. In particular, we do not know what substrates have to be phosphorylated for commitment to occur (although in mammalian cells, Rb is almost certainly one of the substrates).

The purpose of the present work is to develop methods for identifying substrates of the cyclin-Cdk complexes. In particular, we are interested in G1 substrates. To begin, experiments will be done in the yeast *S. cerevisiae*, and then the project will expand into mammalian cells.

There are two main approaches. The first approach uses two-dimensional gels to examine phosphoproteins. Various cyclins are expressed from a *GAL* promoter, and cells with the over-expressed cyclins are labelled with ^{32}P . The pattern of spots on a 2D gel is then compared between cells expressing and not expressing the cyclin. Extra spots in the cyclin-expressing cells may be substrates.

The second approach is to develop antibodies against phosphoserine followed by proline, and phosphothreonine followed by proline. Such antibodies would recognize proteins phosphorylated by Cdk complexes. Thus, such proteins could be immunoprecipitated and sequenced. This could also be combined with the 2D gels as an enrichment step.

6. Body of the Report.

The work has gone somewhat slowly, because it took some time to find a well-qualified post-doctoral fellow. It was not until January that the post-doctoral fellow arrived. Nevertheless, some significant progress has been made.

A. Visualization of G1 Cdc28 substrates on 2D gels.

i. Optimize labeling, extraction, and gel conditions. Quite a number of experiments of this kind have been done. The 2D gels are quite limited in the amount of protein that can be loaded. We have modified previous labelling/extraction procedures so that we now label fewer cells with the same amount of ^{32}P . Thus, we actually load more ^{32}P on the gels, which increases sensitivity markedly. The reduced protein load also increases resolution. Gel conditions were also examined. We found that our original conditions were essentially optimal for pH 4-8 and pH 3-10 gels. However, non-equilibrium (NEQ) gels were poor. The non-equilibrium gels are now much improved, but still not as good as the equilibrium gels. The non-equilibrium gels are very important, because even the "pH 3-10" gels actually focus proteins only up to about pH 7.6. The non-equilibrium gels are essential to look at more basic proteins. Experiments with the non-equilibrium gels are continuing.

We examined ^{33}P as a label instead of ^{32}P . Although it had many advantages in terms of ease of handling and resolution, its specific activity was of course much lower (because of the longer half-life), and this reduces the sensitivity so much that it is not a practical alternative to ^{32}P for these experiments.

In summary, the labeling, extraction, and gel systems are now about as good

as we can make them for proteins of iso-electric points 7.6 and lower. Some improvement is still required for more basic proteins.

ii. Construct strains with different inducible cyclins. This has been nearly completed. We have *GAL-CLN1*, *GAL-CLN2*, *GAL-CLN3*, *GAL-PCL2* (Pcl2, also known as OrfD, is a cyclin of uncertain biological role which normally activates the Pho85 Cdk), *GAL-CLB5* and *GAL-CLB2*. *GAL-CLB3* is under construction. This gives us essentially a complete set of relevant cyclins.

iii. Compare the substrates phosphorylated by the different induced cyclins. We have made only modest progress on this part of the project. To make a comprehensive comparison, we need good non-equilibrium gels so that we can see the proteins with a high isoelectric point. Since we have not had really satisfying gels, we have not put our main effort into these comparisons. However, some preliminary experiments have been done. It appears (on pH 4-8 and 3-10 gels) that *GAL-CLN1*, *GAL-CLN2*, and *GAL-CLN3* induce phosphorylation of the same spots (three strings of spots). However, *GAL-CLB2* does not induce phosphorylation of any of these spots. At first sight, this is very encouraging, because it argues that these spots are cyclin specific. This may be true. However, the *GAL-CLB2* does not induce phosphorylation of any spots we can see, although we know that Clb2-associated histone H1 kinase activity is dramatically induced. The *GAL-CLB2* seems to make the cells rather sick, and inhibit incorporation of ^{32}P . Thus, the failure to see induced phosphorylation may not be meaningful.

In the near future, we will try *GAL-PCL1*, which will be especially interesting, since on the one hand it is expressed in G1, and so in some sense is a "G1 cyclin", but on the other hand its Cdk is Pho80, not Cdc28.

B. Develop antibodies against phosphoSer-Pro and phosphoThr-Pro.

The phosphorylated peptide antigens are in the process of being synthesized.

In the meantime, it has become apparent that a pre-existing monoclonal antibody called MPM2 may be exactly the kind of reagent we are trying to make (Westendorf et al. 1994). MPM2 was discovered many years ago (Davis et al. 1983) as a monoclonal antibody detecting antigens which appear specifically during mitosis. It then became clear that MPM2 was detecting certain phosphoproteins, and it is now known that the MPM2 epitope includes phosphoSer-Pro (PSP) or phosphoThr-Pro (PTP). Since these are the key residues in a consensus Cdk phosphorylation site (S/T-P-X-K/R), phosphorylated Cdk substrates are amongst MPM2-reactive proteins. Recent work of Westendorf et al. suggests that some very good Cdk sites (e.g., L^{PTPLK}) are excellent MPM2 antigens, while other good sites (e.g., K^{PTPVK}) are poor MPM2 antigens. Thus, MPM2 may react with some substrates but not others. This will probably be true of any other antibody, as well.

Some other MPM2 epitopes are not cdc2 sites, because they do not have the basic residue in the fourth position. Instead, they are probably MAP kinase sites. Nevertheless, if the only requirement for recognition by MPM2 is phospho-Ser-Pro or phosphoThr-Pro, then MPM2 will recognize phosphorylated cdc2/Cdc28 sites as well as MAP kinase sites.

MPM2 has just become commercially available from UBI, and we have recently received our first vial. We look forward to using MPM2 in combination with the *GAL-CLN* ^{32}P -labelling experiment. That is, we will induce a *CLN* or *CLB* cyclin, label with ^{32}P , then immunoprecipitate with MPM2. The control will be uninduced cells. We will then run the immunoprecipitate on 1D and 2D gels, and look for ^{32}P -labelled spots specific to cyclin induction. If such spots are found, we will try to obtain enough for microsequencing, or for identification by other methods as described in the original grant proposal.

In addition, we have prepared some proteins known to be substrates for Cdc28-cyclin complexes (e.g., histone H1, Mcm3, Sic1), and are asking whether MPM2 will recognize these proteins after (but not before) phosphorylation.

C. Other work.

Two possible G1 substrates of Cdc28 are Sic1 and Mcm3. We have obtained purified Sic1 and Mcm3, and have phosphorylated them *in vitro*. Both are excellent substrates. Using 2D gels, we have assayed the number of phosphates added (each phosphate gives one extra spot in the isoelectric focusing dimension). Sic1 can gain up to 12 phosphates (!), whereas Mcm3 seems to get 4 or 5. We have not yet characterized Sic1 *in vivo*, but Mcm3 *in vivo* also seems to get 4 or 5 phosphates.

Personnel paid from the grant were B. Futcher (P.I.), D. Marshak (Co-P.I.), G. Sherlock (postdoc), J.P. Liu (lab technician) and E. Araya (part time lab aide).

7. Conclusions.

Except for the late arrival of the post-doctoral fellow, and consequently a delay in some of the research, the project is proceeding more-or-less as planned. The MPM2 antibody may be a valuable alternative to the anti-³²PSP or ³²TP antibodies originally proposed.

8. References:

1. Davis, F. Tsao, T., Fowler, S. and Rao, P. (1983) PNAS 80, 2926-2930.
2. Westendorf, J.M., Rao, P.N., and Gerace, L. (1994) PNAS 91, 714-718.